Dtsch. Entomol. Z. 72 (2) 2025, 505–516 | DOI 10.3897/dez.72.161108

PENSOFT

MUSEUM FÜR NATURKUNDE BERLIN

Dismegistus madagascariensis sp. nov., the first representative of the family Parastrachiidae (Hemiptera, Heteroptera) in Madagascar

Petr Kment[1], Jerzy A. Lis[2]

1 *Department of Entomology, National Museum of the Czech Republic, Cirkusová 1740, CZ-193 00 Praha 9 – Horní Počernice, Czech Republic*

2 *Institute of Biology, University of Opole, Oleska 22, 45-052 Opole, Poland*

https://zoobank.org/70F64B13-DE49-485F-8FCE-E984CBC0D10B

Corresponding author: Petr Kment (sigara@post.cz)

Academic editor: Viktor Hartung ♦ **Received** 4 June 2025 ♦ **Accepted** 29 October 2025 ♦ **Published** 3 December 2025

Abstract

The species *Dismegistus madagascariensis* **sp. nov.** is described from Andohahela National Park, southeastern Madagascar. It represents the first record of the genus *Dismegistus* Amyot & Serville, 1843, and the family Parastrachiidae (Hemiptera, Heteroptera, Pentatomomorpha, Pentatomoidea) from the island. The new species is compared with the six previously described congeners, and a checklist with their distributions is provided.

Key Words

Afrotropical Region, new record, new species, Pentatomomorpha, Pentatomoidea, taxonomy

Introduction

Since the publication of the Linnean "Systema Naturae" (10th edition, Linnaeus 1758), animal species are not only named by a binominal name (binomen), but also classified in a hierarchical system. While the systematic placement of many taxa did not pose a great challenge to the zoologists of the past, there are some taxa that keep travelling throughout the accepted system. Among the true bugs (Hemiptera, Heteroptera), we can mention several such travellers, for example, *Joppeicus paradoxus* Puton, 1881 (Cimicomorpha, Joppeicidae) (for review, see Roca-Cusachs and Kment 2022), *Meschia* Distant, 1910, and *Heissothignus* Slater & Brailovsky, 2006 (Pentatomomorpha, Meschiidae) (see Malipatil 2014; Malipatil et al. 2021; Lu et al. 2025), *Thaumastella* Horváth, 1896 (Pentatomomorpha, Thaumastellidae) (see Kment et al. 2024), or *Eumenotes* Westwood, 1846 (Pentatomomorpha, Dinidoridae) (see Lis et al. 2012; Kment and Kocorek 2014). In the case of *Serbana borneensis* Distant, 1906 (Pentatomomorpha, Phloeidae or Pentatomidae), the systematic placement remains unresolved even at the beginning of the era of DNA-based phylogenetics and phylogenomics (see Grazia et al. 2008; Rider et al. 2018; Roca-Cusachs et al. 2022; López and Schwertner 2023; Luo et al. 2025). Another genus that has travelled considerably throughout the superfamily Pentatomoidea is *Dismegistus* Amyot & Serville, 1843.

The genus was initially described by Amyot and Serville (1843) for *Cydnus circumcinctus* Hahn, 1834, within the category "Séhirides" (which can be compared to Sehirinae in our current classification) as part of their expansive concept of the "Spinipedes." At that time, the latter group encompassed species from the Cydnidae as well as certain representatives of the existing tribe Sciocorini (Pentatomidae). Dallas (1851) categorized *Dismegistus* alongside *Sehirus* within the family Cydnidae; similarly, Stål (1876) positioned *Dismegistus* close to the sehirine genera within his subfamily "Cydnina." Nonetheless, Stål (1876) considered his "Cydnina" to be a subfamily within the family Pentatomidae rather than a distinct family. In his world monograph of the genera of the Cydnidae (termed "Cydnides"), Signoret (1881–1884) regarded this group at the subfamilial rank within the family Pentatomidae. Despite his extensive interpretation of the Sehirinae (his "Séhirides"), he had already decided (Signoret 1880)

Copyright *Petr Kment & Jerzy A. Lis*. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

to transfer *Dismegistus* to the tribe Strachiini (referred to as "Strachiides"), thereby formally placing it within the family Pentatomidae. Therefore, this genus was not even mentioned in his revision of the world Cydnidae (Signoret 1881–1884).

The majority of subsequent authors followed Signoret (1880) in considering the genus *Dismegistus* a member of Pentatomidae. However, Bergroth (1923) transferred it from the Pentatominae to the Asopinae based on the morphology of the bucculae and labium. In contrast, Hesse (1925), like Jeannel (1913) previously, retained it in the Pentatominae.

Leston (1956) examined the morphology and anatomy of *Dismegistus* in detail, demonstrating that the genus should be excluded from the Pentatomidae and returned to the Cydnidae. He also indicated its undeniable affinity with the Sehirinae and placed it in this subfamily.

In the analysis of the higher classification of the Cydnidae by Dolling (1981), the systematic position of *Dismegistus* was reviewed. Despite his broad conception of the Cydnidae, in which he included also the families Thaumastellidae and Thyreocoridae, Dolling disagreed with Leston (1956), thereby excluding *Dismegistus* from the Cydnidae and categorizing the genus as Pentatomoidea *incertae sedis*.

In the same year, Schaefer (1981) studied the stridulitrum and plectrum in certain "primitive" Pentatomoidea, incorporating a single species of this genus into his investigations, specifically *D.* cf. *fimbriatus* (Thunberg, 1783). However, he was unaware of Dolling's (1981) study, as both papers were published in the same month. Therefore, contrary to Dolling, he placed *Dismegistus* within the subfamily Sehirinae, consistent with the previous classification by Leston (1956).

Following this, Schaefer et al. (1988) aimed to ascertain the correct systematic position of the genus *Parastrachia* Oshanin, 1922, within Pentatomoidea by referencing certain morphological characters of *Dismegistus*. Although the authors cited Bergroth's (1923) paper, they failed to comment on the close relationship between *Dismegistus* and *Parastrachia*, despite Bergroth (1923) having discussed these two genera together.

For over a decade, species of this genus were excluded from analyses of the superfamily Pentatomoidea. This is best illustrated by the fact that even in the most significant work of that period (Schuh and Slater 1995), the genus was not mentioned, despite its previously contentious nature regarding its systematic position within the Pentatomoidea.

Only at the beginning of the 21st century were several papers published describing various morphological characters within the family Cydnidae (Lis and Heyna 2001; Lis and Hohol-Kilinkiewicz 2002a, 2002b; Lis and Schaefer 2005). These included two species of *Dismegistus*, namely *D.* cf. *binotatus* (Westwood, 1837) and *D.* cf. *fimbriatus*. However, in all these studies, the authors interpreted *Dismegistus* as a genus of uncertain systematic position. This treatment was also adopted by Pluot-Sigwalt and Lis (2008) when analyzing the morphology of the Cydnidae spermatheca. Nevertheless, they revealed significant similarities in the structure of this organ in *Dismegistus* and *Parastrachia*.

However, it was not until Grazia et al. (2008) analyzed families within the Pentatomoidea based on morphological features and DNA sequences that the monophyly of a group comprising species representing *Parastrachia* and *Dismegistus* was unequivocally demonstrated. Consequently, they broadened the concept of Sweet and Schaefer (2002) by including *Dismegistus* for the first time in the Parastrachiidae. Subsequent morphological (Lis 2010a, 2010b) and molecular analyses (Wu et al. 2016; Lis and Domagała 2024; Lis et al. 2012, 2024; Luo et al. 2025) have confirmed the affiliation of these two genera to a single monophyletic group, the family Parastrachiidae. However, in the context of nuclear ribosomal DNA gene analyses (Lis et al. 2017), species from both genera did not form a monophyletic group; yet, they were classified within the monophyletic subfamily Sehirinae.

As one can conclude from all the aforementioned data, the systematic position of *Dismegistus* varied not only depending on the author's concept of the families within the Pentatomoidea but also on the characters considered. It was classified either under the Sehirinae or the Cydninae of the Cydnidae, the Pentatominae (Strachiini) or Asopinae of the Pentatomidae, or finally, in a separate family, Parastrachiidae (see also Lis 2025).

The genus *Dismegistus* (from Greek *dis mégistos*, meaning "twice as big") currently includes six valid species, distributed mostly in eastern and southern Africa (Jeannel 1913; Hesse 1925; for a review of the species, see Table 1). The genus was last revised by Jeannel (1913), who provided a key for the five species he considered valid. A catalogue of the genus attempted by Robertson (2009) was incomplete. In this contribution, we describe an additional species of *Dismegistus* from Madagascar, which represents the first record of the genus and the entire family Parastrachiidae from that island (cf. Cachan 1952). The genus is also unknown in the nearby Comoro Islands, sharing a significant part of its fauna with Madagascar (Nicholas et al. 2024). In addition, we review the synonymy and distribution of all six previously described species (see Table 1).

Material and methods

Photographs were taken using a Canon MP-E 65 mm macro lens attached to a Canon EOS 550D camera. Final images were stacked from multiple layers using Helicon Focus 5.1 Pro software. Uncoated specimens were examined with a Hitachi S-3700N environmental scanning electron microscope at the Department of Palaeontology, National Museum of the Czech Republic, Prague. The images were edited using Adobe Photoshop CS (version 8.0).

Table 1. Checklist and distribution of the species of *Dismegistus* Amyot & Serville, 1843, based on original descriptions and published records. The identity of the species and distributional records requires revision.

Species of the genus *Dismegistus*	Distribution
D. binotatus (Westwood, 1837: 8, 34–35, as *Pentatoma binotata*) (synonymy by Stål 1866: 252) = *Cimex sanguineus* DeGeer, 1778: 616, pl. 46: fig. 1 (primary junior homonym of *Cimex sanguineus* Schrank, 1776, Reduviidae) = *Cydnus sanguinolentus* Germar, 1838: 184–185 (synonymy by Dallas 1851: 130) = *Cydnus sanguinolentus* Herrich-Schäffer, 1851: 346, pl. CCCXXIII: fig. 1005 [not 1004] (synonymy by Stål 1865: 28)	Kenya (Jeannel 1913, as *D. sanguineus*), Mozambique (Stål 1865, as *D. binotatus*; 1876, as *D. sanguineus*), Rwanda (Leston 1952; Schouteden 1957), South Africa (Westwood 1837; Dallas 1851; Walker 1867a; Stål 1865, 1876; Distant 1898, both as *D. sanguineus*), Tanzania (Schouteden 1910; Jeannel 1913, both as *D. sanguineus*; Leston 1952)
D. costalis Reiche & Fairmaire, 1849: 438–440 = *Cydnus abyssinicus* Herrich-Schäffer, 1851: 345, pl. CCCXXIII: fig. 1004 [not 1005] (synonymy by Stål 1865: 28) = *Aspongopus rufomarginatus* DeGeer, 1895: 110–111 (synonymy by Schouteden 1905: 15)	Ethiopia (Herrich-Schäffer 1851, as *Cydnus abyssinicus*; Stål 1865, 1876; Lethierry 1881, 1883; Carlini 1895, as *Aspongopus rufomarginatus*; Courteaux 1922; Mancini 1939, 1956), Erithrea (Mancini 1954), Senegal (Reiche and Fairmaire 1849), South Africa (Reiche and Fairmaire 1849)
D. fimbriatus (Thunberg, 1783: 47–48, as *Cimex fimbriatus*) = *Cimex capensis* Gmelin, 1790: 2158 (unnecessary new name for *Cimex fimbriatus* Thunberg, 1783) = *Cydonus* [*sic!*] *circumcinctus* Hahn, 1834: 115, pl. LXV: fig. 195 (synonymy by Germar 1838: 184 and Stål 1865: 29) = *Strachia aenescens* Walker, 1867b: 325–326 (synonymy by Distant 1898: 316)	Ethiopia (Lethierry 1881; Jeannel 1913; Courteaux 1922), Guinea (Villiers 1952), Mozambique (Montandon 1899), Namibia (Schumacher 1913; Hesse 1925), Republic of the Congo (Göllner-Scheiding 2012), South Africa (e.g., Thunberg 1783; Hahn 1834, as *Cydonus circumcinctus*; Dallas 1851; Walker 1867a; Stål 1865, 1876; Wallengren 1875; Distant 1892, 1898; Walker 1867b, as *S. aenescens*), Tanzania (Jeannel 1913), Zimbabwe (Distant 1898, as Mashonaland)
D. funebris Distant, 1900: 169–170, pl. II: fig. 8	Democratic Republic of the Congo (Haut-Katanga) (Schouteden 1913), Tanzania (Distant 1900; Schouteden 1910; Jeannel 1913)
***D. madagascariensis* sp. nov.**	Madagascar (this paper)
D. royeri Jeannel, 1913: 88, 90–91, pl. III: fig. 34	Angola, Kenya, South Africa (Natal) (Jeannel 1913)
D. rufomarginatus Hesse, 1925: 35, pl. 1: fig. 5	Namibia (Hesse 1925)

The following dimensions were measured: body length (from the apex of the mandibular plates to the apex of the abdomen, dorsal view); head length (from the apex of the mandibular plates to the anterior margin of the pronotum, frontal view); head width (width of the head including compound eyes); interocular width (between inner margins of compound eyes); length of each antennomere and labiomere; pronotum length (medially, from the anterior to the posterior margin of the pronotum in anterodorsal view); pronotum width (maximum width between the humeri in dorsal view); scutellum length (medially from the base to the apex); and scutellum width (maximum width at the base between the basal angles of the scutellum).

The morphological terms used follow Tsai et al. (2011), Rédei and Tsai (2021, 2022), Roca-Cusachs et al. (2024), and Salini et al. (2023, 2025). The specialized terms derive from the following papers: antennomeres – Zrzavý (1990); external scent efferent system of the metathoracic scent glands – Kment and Vilímová (2010); trichobothria – Gao et al. (2017); female external genitalia – Zhou and Rédei (2020).

In the examined holotype, the data on labels are quoted verbatim, with a slash (/) separating data on different rows of a single label and a double slash (//) dividing data on different labels. The authors' comments are given in square brackets [], and the following abbreviations are used: [hw] = handwritten, [p] = printed. Unless stated otherwise, the described labels are off white.

Specimens deposited in the following institutional collections were examined or mentioned in this study:

MMBC Moravian Museum, Brno, Czech Republic;
MFNB Museum für Naturkunde, Berlin, Germany;
MNHN Muséum National d'Histoire Naturelle, Paris, France;
MRAC Royal Museum for Central Africa, Tervuren, Belgium;
NMPC National Museum of the Czech Republic, Prague, Czech Republic.

We used the following specimens for direct comparison with the holotype of *D. madagascariensis* sp. nov. Due to the absence of a modern revision of the genus, the species identifications could only be tentative (cf. Lis and Domagała 2024).

Dismegistus* cf. *binotatus (Westwood, 1837): **KENYA:** Nairobi, 1 ♀, Machul[ka lgt.] (NMPC). Ikutha, 1 ♀, x/xii.1960, Muche lgt., ex col. Breuning, Coll. Mus. Tervuren (MRAC). **SOUTH AFRICA:** Mpumalanga, Waterval Boven, Elands river, 10.–11.ii.2000, 1 ♂ [Fig. 4], S. Bečvář lgt. (MMBC). Transvaal, Rustenburg, 9.xii.1950, A. L. Capener lgt., 1 ♂ 2 ♀♀, R. I. Sailer det. as *D. binotatus* (MMBC). **TANZANIA:** T.T.O-Afrika, Marangu, 1.–20.iii.1959, 1 ♂ 2 ♀♀, Lindner leg. (NMPC).

Dismegistus* cf. *fimbriatus (Thunberg, 1783): **SOUTH AFRICA:** Bontebok Nat. Park, 16.–18.xi.1993, 1 ♀, F. Koch lgt. (MFNB). Bothaville, Orange Fr. St., 22.xii.1998, 1 ♂, 15.i.1999, 1 ♀, Dr. Brauns lgt. (MRAC). Cape of Good Hope N.R., 34°16'S, 18°23'E, 24.–25.iv.1995, 1 ♂ 1 ♀, J. Deckert lgt. (MFNB). Cape Province, De Hoop Nature Res., 16.–18.xi.1993, 34°27'S, 20°24'E, 1 ♂, J. Deckert lgt. (MFNB). 'Holub', 2 ♂♂ 5 ♀♀ (NMPC; for collecting details see Kment and Rédei 2018). Mpumalamga, Phabeni, 14.i.2007, 1 ♂, S. Snäll lgt. (NMPC). Mpumalanga, Waterval Boven, Elands river, 10.–11.ii.2000, 2 ♂♂ 1 ♀ [Fig. 5], S. Bečvář lgt. (MMBC). **ZIMBABWE:** Rhodesia mer., Antelope, hospital, i.–ii.1972, 2 ♂♂, Z. Cakl lgt. (MMBC).

Dismegistus royeri Jeannel, 1913: **SOUTH AFRICA:** Natal, 1 spec. (syntype) [Figs 6–7], Dr. Martin lgt., R. Jeannel det. (MNHN).

Results

Dismegistus madagascariensis sp. nov.

https://zoobank.org/72BE62AA-4E0C-4505-B351-6866BFC6A7C5

Figs 1–3, 8–19

? *Dismegistus pauliani* (nomen nudum): Leston (1956): 92 (Madagascar).

Type locality. South-east Madagascar, Anosy mountain range, Andohahelo plateau [Andohahela National Park], SW of Trafonaomby Mt., 1770–1950 m a.s.l. (ca. 24°33'54"S, 46°43'14"E).

Type material. ***Holotype*:** ♀ (Figs 1–3), 'Chaînes anosyennes / S. O. du Trafonaomby / plateau Andohahelo / 1770· 1950 m, V. - 1972 [p] // Museum Paris [p] / A. Peyrieras / I. 1974 [hw] // ♀ [p] // HOLOTYPUS / *DISMEGISTUS* / *MADAGASCARIENSIS* / sp. nov. / des. P.Kment & J.A.Lis 2025 [p, red label]' (MNHN). The holotype was remounted on a new piece of card, with a trace of past pinning on the scutellum; hemelytra slightly spread, abdominal ventrites depressed (Fig. 2); both distiflagella, left basiflagellum, both protarsi, right mesotarsus and left metatibia and metatarsus missing (Fig. 1).

Description (female). Coloration (Figs 1, 2). Body, antennae, and legs black, pronotum and propleura bright orange-red; apex of scutellum with narrow reddish-brown margin; clavi black; coria (including hypocostal lamina) reddish-black, basally and along each R+M vein darker (against the surface of abdomen corium appearing black), lateral margins of coria dark red; membranes translucent, fumose brown; anterolateral angle of each mesepisternum red, metepimera reddish-black (Fig. 2).

Structure. Head triangular in dorsal (i.e., frontal) view (Fig. 10), declivous (Figs 8, 9). Mandibular plates slightly surpassing apex of clypeus to form anterior notch (Figs 10, 11). Lateral margins of head elevated dorsally in form of ridges, remaining surface of mandibular plates and clypeus flat (Fig. 10), vertex between eyes and ocelli convex (Fig. 9). Bucculae crescent shaped, ca. as long as labiomere I (Fig. 9). Labium (Fig. 14) reaching middle of mesosternum (Fig. 15). Length of labiomeres III > IV > II > I, labiomere I reaching about head midlength (Fig. 14). Antenna with scape (I) shortest, stoutest, basipedicellite (IIa) and distipedicellite (IIb) about same length, cylindrical, only slightly wider apically than at base (Figs 1, 2), basiflagellum (III) spindle-shaped, slightly flattened, longer than preceding antennomeres, distiflagellum (IV) missing in examined specimen.

Pronotum (Figs 1, 12) with anterior margin shallowly concave, anterolateral and humeral angles broadly rounded; lateral margins narrowly explanate, convex in dorsal view, slightly curved dorsally in lateral view; posterior margin slightly concave in middle. Pronotal disc with very shallow, almost indistinct sublateral depression anterolaterally (Fig. 12). Scutellum triangular with convex surface, insinuate at frenal apices, apex of scutellum broadly rounded posteriorly (Figs 1, 13).

Coria somewhat explanate laterally each with well developed hypocostal lamina (Fig. 2); costal margins of coria convex in dorsal view (Figs 1, 2, 13), each exocorium with depressed medial furrow (Fig. 13: mf) anteriorly between Sc and R+M veins, posterior margin of each corium rounded (Fig. 1); veins of membranes translucent, indistinct.

Thoracic sterna neither grooved or keeled (Fig. 15). External scent efferent system of metathoracic scent glands (Fig. 16) reduced, ostioles V-shaped (Fig. 16: o), each situated between meso- and metacoxal cavities, vestibulum and vestibular scars very short, peritremes not emarginated, peritremal sufaces merging with surrounding metapleura, only median furrow well-developed (Fig. 16: mf); evaporatoria (Fig. 16: ev) rudimentary, mycoid surface limited to small area posteriad of ostioles, elongate patch along suture between meso- and metapleura. Femora simple, oval in cross-section; tibiae with dorsal surface shallowly grooved (Fig. 8). Tarsomere I stoutest, longest, longer than II + III combined, tarsomere III longer than II.

Each abdominal ventrite bearing two trichobothria postero-/posterolaterad of each spiracle (Fig. 18: sp, tr), bothrium of type A (Fig. 19). Female terminalia (Fig. 17) with valvifers VIII semicircular with posterior margin slightly rounded (Fig. 17: vf8), laterotergites IX small, triangular, with lateral margins rounded (Fig. 17: lt9); laterotergites VIII medially fused (Fig. 17: lt8).

Integument and vestiture. Pronotum and head shining, scutellum and hemelytra submatte (Fig. 1). Head (Figs 10, 11) impunctate, clypeus basally and apically with shallow transverse wrinkles, mandibular plates laterally with transverse wrinkles fading centrally (Fig. 10). Pronotum with sparse, shallow, concolorous punctures, more pronounced in center of disc (Fig. 12). Scutellar disc with concolorous punctures, sparse, shallow basally, more pronounced, connecting with transverse wrinkles in central and posterior parts of disc (Fig. 13). Clavi and coria with dense concolorous punctures, deeper, denser than on pronotum. Ventral surface of body nearly smooth (Figs 14–19), metepimera with large shallow punctures.

Body bare except distipedicellite (IIb) and basiflagellum (III) with fine, short, semierect pilosity. Each tibia with four rows of stout black semierect spines, one row dorsally on each side of of flattened dorsal surface (Fig. 8), two rows on ventral surface, their spines forming acute angle ventrally; besides the stout spines each tibia bearing additional finer, shorter setae. Tarsi with short, semierect to erect setae, especially dense on ventral surfaces.

Measurements (mm). Body length 8.33; head: length 1.35, width (including compound eyes) 2.10, interocular width 1.37; lengths of labiomeres: I – 0.35, II – 0.60, III – 0.82, IV – 0.76; lengths of antennomeres: I – 0.52, IIa – 1.02, IIb – 1.00, III – 1.32, IV – missing; pronotum: length 2.42, width 5.05; scutellum: length 3.23, width 2.94.

Male. Unknown.

Figures 1–7. Habitus of *Dismegistus* species. **1–3.** *Dismegistus madagascariensis* sp. nov., ♀, holotype (**1.** Dorsal view, **2.** Ventral view, **3.** Holotype labels; body length: 8.33 mm); **4.** *D.* cf. *binotatus* (Westwood, 1837), ♂, South Africa: Waterval Boven, dorsal view (body length: 9.31 mm); **5.** *D.* cf. *fimbriatus* (Thunberg, 1783), ♀, South Africa: Waterval Boven, dorsal view (body length: 8.63 mm); **6, 7.** *D. royeri* Jeannel, 1913, ♀, syntype, South Africa: Natal: **6.** Dorsal view, **7.** Type labels. Scale bar: 1 mm.

Differential diagnosis. The new species is similar in body length to *D.* cf. *fimbriatus*, *D.* cf. *funebris* Distant, 1900, and *D. royeri* Jeannel, 1913. The females of these species, measuring approximately 5.0 mm to nearly 9.0 mm in length, are smaller than those of other species within this genus, such as *D.* cf. *binotatus*, *D.* cf. *costalis* Reiche & Fairmaire, 1849, and *D.* cf. *rufomarginatus* Hesse, 1925, where the female body length typically









Figures 8–13. Morphology of *Dismegistus madagascariensis* sp. nov., ♀, holotype. **8.** Head, pro- and metathorax in lateral view (magnification 32×); **9–11.** Head: **9.** Lateral view (65×); **10.** Frontal view (60×); **11.** Ventral view (60×); **12.** Pronotum (anterodorsal view, 25×); **13.** Scutellum, clavus, and anterior portion of corium (23×). Lettering: mf – medial furrow, R+M – R+M vein. Scale bars: 0.5 mm (**9–11**); 1 mm (**8, 12, 13**).

exceeds 10.0 mm and can occasionally reach as much as 16.0 mm. However, *D. madagascariensis* can be easily distinguished from *D.* cf. *fimbriatus*, *D.* cf. *funebris*, and *D. royeri* by the coloration of the pronotum and hemelytra. The pronotum of *D. madagascariensis* is uniformly red on the dorsal side, unlike the other three species, where only the lateral portions and posterior parts of the pronotum show red. The same applies to the outer sections of the exocoria, which in *D.* cf. *fimbriatus*, *D.* cf. *funebris*, and *D. royeri* are predominantly red along their length and distinctly differ in color from the darker mesocoria. In *D. madagascariensis*, the base of each exocorium is noticeably darker, resembling the color of the mesocorium. Furthermore, unlike all other species in the genus, the hemelytra of *D. madagascariensis* are broadly expanded laterally, reaching distinctly laterad of the humeral angles.

Etymology. The species epithet is the Latinized adjective *madagascariensis* (*-is*, *-e*), referring to the area of the species distribution.

Habitat and biology. Unknown. The type locality, Andohahela National Park, is situated at the southern end of the Malagasy Highlands. The park is divided into three













Figures 14–19. Morphology of *Dismegistus madagascariensis* sp. nov., ♀, holotype. **14.** Labium (magnification 55×); **15.** Meso- and metasternum (40×); **16.** Ostiole and evaporatorium of metathoracic scent glands (170×); **17.** Female terminalia (70×); **18.** Spiracle and trichobothria on ventrite III (170×); **19.** Detail of trichobothria on ventrite III (500×). Lettering: ev – evaporatorium, la – labiomere I, lr – labrum, lt8–9 – laterotergites VIII–IX, mf – median furrow of peritreme, o – ostiole of metathoracic scent gland, ms – mesosternum, mt – metasternum, sp – spiracle, tr – trichobothrium, vf8 – valvifer VIII. Scale bars: 0.1 mm (**19**); 0.2 mm (**16, 18**); 0.5 mm (**14, 15, 17**).

zones. The first zone, Malio, ranges from 100 meters to the summit of Pic d'Andohahela at 1,956 meters, and has dense lowland and montane rainforest. The second zone, Ihazofotsy-Mangatsiaka, contains dry spiny forest at altitudes ranging from 100 to 1,005 meters at the summit of Pic de Vohidagoro. The third zone, Tsimelahy, is mainly at an altitude of 125 meters and contains the unique Ranopiso transitional forest. The mountains form a natural barrier to the moist trade winds that blow from the east, causing on the eastern side a rainfall of 1,500–2,000 millimeters per year that supports one of the few rain forests south of the Tropic of Capricorn. At the park's western edge, the rainfall is just

600–700 millimeters per year, and the resulting vegetation is a dry, spiny forest characteristic of southern Madagascar (Paulian et al. 1973; Anonymus 2025). As the holotype label indicates, the specimen was collected at the leeward south-west slopes of Trafonaomby Mt. (1770–1950 m a.s.l.), i.e., just under the mountain top. This suggests its finding in the forest zone, though the exact type of forest would be just a speculation. Collecting fresh specimens is necessary to elucidate the bionomy and habitat requirements of the new species. The available habitat information concerning the species in continental Africa concerns relatively drier habitats (e.g., Lis and Domagała 2024).

Distribution. Known only from the type locality, Trafonaomby Mt. in south-east Madagascar.

Remark. When examining species of *Dismegistus*, Leston (1956) noted that his study was based on material from various regions of South Africa (*D.* cf. *fimbriatus*, *D.* cf. *binotatus*) and Madagascar (*D. pauliani*). Although a Latin name was assigned to the species from Madagascar, Leston classified it as a *nomen nudum*. Concurrently, he suggested that the species would be described once sufficient material became available to undertake a comprehensive revision of the entire genus *Dismegistus*. Unfortunately, this species from Madagascar was neither described by Leston nor mentioned in his subsequent publications. Consequently, it is currently difficult to determine whether the specimens used as the basis for *D. pauliani* represent the same species described in this paper. Nevertheless, given that these are the only specimens of this genus recorded from Madagascar, the name *D. pauliani* has been tentatively included as a *nomen nudum* under *D. madagascariensis*.

Discussion

Jeannel (1913) distinguished the species of *Dismegistus* based on the size, shape, and coloration of the body, the lengths of the antennomeres, and the structure of the pronotum. Hesse (1925) also used the same characters to distinguish the sixth species he described, *Dismegistus rufomarginatus*. Some general morphological characters were described and illustrated by Leston (1956) and Dolling (1981), including the details of the male (paramere, penis) and female genitalia (terminalia, spermatheca) of *D.* cf. *binotatus* and *D.* cf. *fimbriatus*—the latter (female terminalia) not much different from the condition in *D. madagascariensis* (Fig. 17). These structures must be analyzed and described in detail in the future, as they differ only slightly across all species (J. A. Lis, unpubl. data). This will form the basis for further analyses of all *Dismegistus* species and a comprehensive revision of the genus.

Due to its unique history, Madagascar is considered one of the most significant hotspots of biodiversity (e.g., de Wit 2003; Ganzhorn et al. 2008; Ralimanana et al. 2022; Jimenez et al. 2023), and many new species of insects are described from Madagascar every year, including many true bugs (Heteroptera) (e.g., Guilbert 2020; Kim et al. 2020; Kondorosy et al. 2020; Zettel 2020; Zettel and Laciny 2020; Matocq 2021; Chen et al. 2022; Sites and Bergsten 2022; Taszakowski et al. 2022, 2025a, b; Zámbó et al. 2022; Kóbor 2023; Masłowski et al. 2023; Gierlasiński et al. 2024; Heiss and Eckelt 2024; Chen and Cai 2025; Cook 2025; this paper). These discoveries mostly represent new genera and species belonging to family-group taxa previously known to occur in Madagascar. However, some of the endemic taxa described in the past three decades also represent discoveries of tribes, subfamilies, or even families previously unknown in Madagascar. Brailovsky (2007) described *Madagalesus* Brailovsky, 2007, a new genus including two new species and representing the first record of the tribe Dasynini (Coreidae, Coreinae). Brailovsky (2011) also added Latimbini, another tribe of Coreinae, based on four new Malagasy species of the genus *Latimbus* Stål, 1859. O'Donnell and Schaefer (2011) first recorded the tribe Lilliputocorini (Rhyparochromidae) based on five undescribed species of the genus *Lilliputocoris* Slater & Woodward, 1979 (see also Kment et al. 2016). Heiss and Eckelt (2024) described the first endemic genus and species of the flat bug subfamily Calisiinae (Aradidae), *Microcalisius madagascariensis* Heiss & Eckelt, 2024. Gierlasiński et al. (2024) described *Kojderus* Magnien & Gierlasiński, 2024, a new monotypic species of Tessaratomidae, representing the first record of the subfamily Natalicolinae from Madagascar. Taszakowski et al. (2025b) described *Psallops madagascariensis* Taszakowski, Kim & Gierlasiński, 2025, the first Malagasy representative of the subfamily Psallopinae (Miridae). Finally, we are aware of additional new taxa still awaiting formal description, such as a new species of the genus *Aeptus* Dallas, 1851 from the tribe Aeptini (Pentatomidae, Pentatominae) (Rider et al. 2018) and the existence of undescribed new taxa of Urostylididae (P. Štys, pers. comm. in Zhou and Rédei 2018; P. Kment, pers. observ.). In this context, the discovery of the first representative of the family Parastrachiidae presented in this paper is not a major surprise.

Acknowledgments

We are obliged to the late Professor Pavel Štys (Faculty of Science, Charles University, Prague, Czech Republic), who first noticed the here described specimen in MNHN, marked it as an undescribed species, and borrowed it for description. We further thank Petr Baňař (Moravian Museum, Brno, Czech Republic) for providing access to P. Štys's collection; Éric Guilbert and Dominique Pluot-Sigwalt (MNHN) for their kind assistance during the first author's visit to MNHN; and David A. Rider (Fargo, North Dakota, USA). Philippe Magnien and D. Pluot-Sigwalt (MNHN) kindly helped us with the translation of the type label in French. We also extend our gratitude to all collection curators at the institutions mentioned in the "Material and methods" section for loaning specimens for our study. The work of P. Kment was financially supported by the Ministry of Culture of the Czech Republic (DKRVO 2024–2028/5.I.b). Museum für Naturkunde (Berlin) waived the publication costs of this article.

References

Amyot CJB, Serville JGA (1843) Histoire naturelle des insectes Hemipteres. Librairie Encyclopedique de Roret ed., Paris, 1 xxvi+ 675 pp. https://doi.org/10.5962/bhl.title.8471

Anonymus (2025) Andohahela National Park. Wikipedia. https://en.wikipedia.org/wiki/Andohahela_National_Park [accessed 2 January 2025]

Bergroth E (1923) On the systematic position of the genera *Dismegistus* Am. S. and *Parastrachia* Dist. Annales de la Société Entomologique de Belgique 63: 70–72.

Brailovsky H (2007) A new genus and two new species of Dasynini (Hemiptera: Heteroptera: Coreidae) from Madagascar. Proceedings of the Entomological Society of Washington 109: 847–855.

Brailovsky H (2011) Insecta, Hemiptera, Heteroptera, Coreidae. Faune de Madagascar. Vol. 94. IRD Éditions, Éditions Quae and Muséum National d'Histoire naturelle, Paris et Marseille, 275 pp.

Cachan P (1952) Les Pentatomidae de Madagascar (Hemipteres Heteropteres). Mémoires de l'Institut Scientifique de Madagascar. Série E, Entomologie 1: 231–462.

Chen Z, Cai W-Z (2025) The second Malagasy species of *Anacanthiocnemis* (Hemiptera: Heteroptera: Reduviidae) with an updated key to the species of the genus. Acta Entomologica Musei Nationalis Pragae 65(1): 21–30. https://doi.org/10.37520/aemnp.2025.003

Chen Z, Li H, Cai W-Z (2022) A new species of the genus *Monicacoris* Putshkov (Hemiptera: Reduviidae: Emesinae) from Madagascar. Annales de la Société Entomologique de France 58(2): 165–172. https://doi.org/10.1080/00379271.2022.2054468 [Nouvelle Serie]

Cook JL (2025) *Paraplea* Esaki and China (Hemiptera: Heteroptera: Pleidae) of Madagascar, a new species and redescription of *Paraplea hovana* (Kirkaldy, 1899). Proceedings of the Entomological Society of Washington 126(4) [2024]: 441–453. https://doi.org/10.4289/0013-8797.126.4.441

Courteaux R (1922) Hémiptères Hétéroptères. In: Voyage de M. le Baron Maurice de Rothschild en Éthiopie et en Afrique Orientale Anglaise (1904–1905). Résultats scientifiques. Animaux Articulés. Imprimerie Nationale, Paris, 263–293.

Dallas WS (1851) List of the specimens of hemipterous insects in the collection of the British Museum. Part 1. Trustees of the British Museum, London, 1–368. [pls. 1–11]

de Carlini A (1895) VII. Rincoti. In: Esplorazione del Giuba e dei suoi affluenti compiuta dal Cap. V. Bottego durante gli anni 1892–93 sotto gli auspicii della Società Geografica Italiana. - Risultati zoologici. Annali del Museo Civico di Storia Naturale di Genova, Serie 2 15: 105–125.

de Wit MJ (2003) Madagascar: Heads it's a continent, tails it's an island. Annual Review of Earth and Planetary Sciences 31(1): 213–248. https://doi.org/10.1146/annurev.earth.31.100901.141337

DeGeer C (1778) Mémoires pour servir a l'histoire des insectes. Tome septieme. Pierre Hesselberg, Stockholm, 950 pp. [+ pls 1–49]

Distant WL (1892) A naturalist in the Transvaal. R. H. Porter, London, xvi + 277 pp. [+ pls I–V] https://doi.org/10.5962/bhl.title.2478

Distant WL (1898) Rhynchota from the Transvaal, Mashonaland, and British Nyasaland. Annals & Magazine of Natural History Series 7 2(10): 294–316. https://doi.org/10.1080/00222939808678047

Distant WL (1900) Undescribed genera and species belonging to the Rhynchotal family Pentatomidæ. Transactions of the Entomological Society of London 1: 163–174. [+ pl. 2] https://doi.org/10.1111/j.1365-2311.1900.tb00998.x

Dolling WR (1981) A rationalized classification of the Burrower Bugs (Cydnidae). Systematic Entomology 6(1): 61–76. https://doi.org/10.1111/j.1365-3113.1981.tb00016.x

Ganzhorn J, Lowry P II, Schatz G, Sommer S (2008) The Biodiversity of Madagascar: One of the world's hottest hotspots on its way out. Oryx 35(4): 346–348. https://doi.org/10.1046/j.1365-3008.2001.00201.x

Gao C-Q, Rédei D, Shi X-Q, Cai B, Liang K, Gao Sh, Bu W-J (2017) A comparative study of the abdominal trichobothria of Trichophora, with emphasis on Lygaeoidea (Hemiptera: Heteroptera). European Journal of Entomology 114: 587–602. https://doi.org/10.14411/eje.2017.072

Germar EF (1838) Hemiptera Heteroptera promontorii Bonæ Spei, nundum descripta, quæ collegit C. F. Drège. Silbermann's Revue Entomologique 5 [1837]: 121–192.

Gierlasiński G, Magnien Ph, Kepel A, Wiśniewski K, Lange A (2024) *Kojderus maximus*, a new genus and species of Tessaratomidae (Hemiptera: Heteroptera), the first Natalicolinae from Madagascar. Acta Entomologica Musei Nationalis Pragae 64(2): 327–335. https://doi.org/10.37520/aemnp.2024.023

Gmelin JF (1790) Caroli a Linné Systema Naturae. Tom I. Pars IV. Beer, Lipsiae, 1517–2224.

Göllner-Scheiding U (2012) Heteroptera aus der Republik Kongo, gesammelt von Dr. G. Hertzel in den Jahren 1977 und 1978. Edessana 2: 95–107.

Grazia J, Schuh RT, Wheeler WC (2008) Phylogenetic relationships of family groups in Pentatomoidea based on morphology and DNA sequences (Insecta: Heteroptera). Cladistics 24(6): 932–976. https://doi.org/10.1111/j.1096-0031.2008.00224.x

Guilbert E (2020) New species of Tingidae (Insecta: Heteroptera) from Madagascar. Zootaxa 4759(3): 391–404. https://doi.org/10.11646/zootaxa.4759.3.5

Hahn CW (1834) Die Wanzenartigen Insekten, getreu nach der Natur abgebildet und beschrieben. Vol. 2. C. H. Zeh'schen Buchhandlung, Nürnberg, 101–120.

Heiss E, Eckelt A (2024) *Microcalisius madagascariensis* gen.n. et sp.n., the first record of Calisiinae (Heteroptera: Aradidae) from Madagascar. Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 76: 1–6.

Herrich-Schäffer GAW (1851) Die Wanzenartigen Insekten. Vol. 9. Lotzbeck, Nürnburg, 257–348.

Hesse AJ (1925) 1. Contributions to a knowledge of the fauna of South-West Africa. IV. A list of the Heteropterous and Hemipterous Hemiptera of South-West Africa. Annals of the South African Museum 23: 1–190. [8 pls]

Jeannel R (1913) I. Pentatomidae. Pp. 1–114 + pls. i–iv. In: Voyage de Ch. Alluaud et R. Jeannel en Afrique orientale, 1911–1912. Résultats scientifiques. Insectes Hémipteres. A. Schulz, Paris, 333 pp. [+ 12 pls]

Jimenez BO, Monatno R, Rothman RS, Williams RC, Wright PC (2023) The biodiversity of an old-growth forest amidst a scorched landscape in Madagascar. Conservation Science and Practice 5(9, e12993): 1–10. https://doi.org/10.1111/csp2.12993

Kim J-G, Roca-Cusachs M, Lim J-G, Jung S-H (2020) *Malagasycoelum dracula* gen. and sp. nov., a new genus and a new species of the subfamily Mirinae (Hemiptera: Heteroptera: Miridae) from Madagascar. Zootaxa 4808(1): 165–170. https://doi.org/10.11646/zootaxa.4808.1.10

Kment P, Kocorek A (2014) *Afromenotes hirsuta*, a new genus and species of Eumenotini from the Democratic Republic of the Congo (Hemiptera: Heteroptera: Dinidoridae). Acta Entomologica Musei Nationalis Pragae 54(1): 109–131.

Kment P, Rédei D (2018) A revision of the types of Heteroptera species described by Géza Horváth based on specimens from collections of Ladislav Duda and Emil Holub. Acta Entomologica Musei Nationalis Pragae 58(1): 275–295. https://doi.org/10.2478/aemnp-2018-0025

Kment P, Vilímová J (2010) Thoracic scent efferent system of the Tessaratomidae sensu lato (Hemiptera: Heteroptera: Pentatomoidea) with implication to the phylogeny of the family. Zootaxa 2363(1): 1–59. https://doi.org/10.11646/zootaxa.2363.1.1

Kment P, Hemala V, Baňař P (2016) *Rhyparoclava pyrrhocoroides*, a new genus and species of autapomorphic Rhyparochromidae with clavate antennae from Madagascar (Hemiptera: Heteroptera). Acta Entomologica Musei Nationalis Pragae 56(2): 517–545.

Kment P, Jacobs DH, Carapezza A, Deckert J, Rider DA, Kóbor P (2024) Desert Bugs (Hemiptera: Heteroptera: Thaumastellidae): New records and review of the distribution and habitat of this relict group. Zootaxa 5541(2): 144–162. https://doi.org/10.11646/zootaxa.5541.2.2

Kóbor P (2023) Big-eyed bugs of the Malagasy Region (Hemiptera: Heteroptera: Lygaeoidea: Geocoridae). Acta Zoologica Academiae Scientiarum Hungaricae 69(3): 247–264. https://doi.org/10.17109/AZH.69.3.247.2023

Kondorosy E, Zámbó A, Kovács S, Baňař P (2020) *Scobinigaster*, a new genus of Lethaeini from Madagascar (Hemiptera: Heteroptera: Rhyparochromidae). Acta Entomologica Musei Nationalis Pragae 60(1): 343–352. https://doi.org/10.37520/aemnp.2020.020

Leston D (1952) Notes on the Ethiopian Pentatomoidea (Hemiptera). – VI. Some insects in the Hope Department, Oxford. Annals & Magazine of Natural History Series 12 5(58): 893–904. https://doi.org/10.1080/00222935208654367

Leston D (1956) The Ethiopian Pentatomoidea (Hemiptera). XXII. On *Dismegistus* Amyot & Serville (Cydnidae). Proceedings of the Royal Entomological Society of London, A 31: 87–94. https://doi.org/10.1111/j.1365-3032.1956.tb00214.x

Lethierry L (1881) Spedizione Italiana nell'Africa Equatoriale. Resultati zoologici. Emitteri. Annali del Museo Civico di Storia Naturale di Genova 16: 277–298.

Lethierry L (1883) Spedizione Italiana nell'Africa Equatoriale. Resultati zoologici. Emitteri. Parte Seconda. Annali del Museo Civico di Storia Naturale di Genova 18: 741–756.

Linnaeus C (1758) Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, diferentiis, synonymis, locis. Editio decima, reformata. Holmiae, 823 + I pp. https://doi.org/10.5962/bhl.title.542

Lis JA (2010a) Coxal combs in the Cydnidae sensu lato and three other related "cydnoid" families – Parastrachiidae, Thaumastellidae, Thyreocoridae (Hemiptera: Heteroptera): functional, taxonomic, and phylogenetic significance. Zootaxa 2476(1): 53–64. https://doi.org/10.11646/zootaxa.2476.1.6

Lis JA (2010b) Pretarsal structures in the family Parastrachiidae (Hemiptera: Heteroptera: Pentatomoidea). Zootaxa 2693(1): 60–62. https://doi.org/10.11646/zootaxa.2693.1.5

Lis JA (2025) Classification of the burrower bugs (Hemiptera: Heteroptera: Cydnidae): A never-ending story? Insects 16(874): 1–15. https://doi.org/10.3390/insects16090874

Lis JA, Domagała PJ (2024) First reports of parental care in species of the genus *Dismegistus* (Hemiptera: Heteroptera: Parastrachiidae). European Journal of Entomology 121: 396–399. https://doi.org/10.14411/eje.2024.043

Lis JA, Heyna J (2001) The metathoracic wing stridulitrum of the Cydnidae (Hemiptera: Heteroptera). Polish Journal of Entomology 70(4): 221–243.

Lis JA, Hohol-Kilinkiewicz A (2002a) Abdominal trichobothrial pattern and its taxonomic and phylogenetic significance in Cydnidae (Hemiptera: Heteroptera). III. Sehirinae. Annales Zoologici (Warszawa) 52: 359–362.

Lis JA, Hohol-Kilinkiewicz A (2002b) Adult dorso-abdominal scent glands in the burrower bugs (Hemiptera: Heteroptera: Cydnidae). Polish Journal of Entomology 71(4): 359–395.

Lis JA, Schaefer CW (2005) Tibial combs in the Cydnidae (Hemiptera: Heteroptera) and their functional, taxonomic and phylogenetic significance. Journal of Zoological Systematics and Evolutionary Research 43(4): 277–283. https://doi.org/10.1111/j.1439-0469.2005.00328.x

Lis JA, Lis P, Ziaja DJ, Kocorek A (2012) Systematic position of Dinidoridae within the superfamily Pentatomoidea (Hemiptera: Heteroptera) revealed by the Bayesian phylogenetic analysis of the mitochondrial 12S and 16S rDNA sequences. Zootaxa 3423(1): 61–68. https://doi.org/10.11646/zootaxa.3423.1.5

Lis JA, Ziaja DJ, Lis B, Gradowska P (2017) Non-monophyly of the "cydnoid" complex within Pentatomoidea (Hemiptera: Heteroptera) revealed by Bayesian phylogenetic analysis of nuclear rDNA sequences. Arthropod Systematics & Phylogeny 75(3): 481–496. https://doi.org/10.3897/asp.75.e31918

Lis JA, Domagała PJ, Lis B (2024) New molecular phylogenetic evidence confirms independent origin of coxal combs in the families of the 'cydnoid' complex (Hemiptera: Heteroptera: Pentatomoidea). Insects 15 (10) (792): 1–15. https://doi.org/10.3390/ijms25020939

López GEL, Schwertner CF (2023) Synopsis of the remarkable family Phloeidae (Hemiptera: Pentatomoidea): species identification, chromatic polymorphism and updated distribution. Journal of Natural History 57(45–48): 2083–2113. https://doi.org/10.1080/00222933.2023.2284417

Lu Y, Ye F, Li J-L, Gao C-Q, Wang Y-H, Shi X-Q, Su Y-Q, Xue H-J, Bu W-J, Zheng L-Y, Xie Q (2025) Phylogenomics of the superfamily Lygaeoidea with proposals on taxonomic system (Hemiptera: Heteroptera). Molecular Phylogenetics and Evolution 209(108356): 1–10. https://doi.org/10.1016/j.ympev.2025.108356

Luo J-Y, Wu Y-Z, Kment P, Salomão AT, Damken C, Wang Y-H, Xie Q (2025) Origin of the only myrmecomorphic stink bug, *Pentamyrmex spinosus* (Hemiptera: Pentatomidae), in the radiation era of ants (Hymenoptera: Formicidae). Systematic Entomology 50(1): 415–427. https://doi.org/10.1111/syen.12664

Malipatil MB (2014) Meschiidae, a new family of Lygaeoidea (Hemiptera: Heteroptera) from India and Australia, with descriptions of a new genus and two new species. Zootaxa 3815(2): 233–248. https://doi.org/10.11646/zootaxa.3815.2.4

Malipatil MB, Yeshwanth HM, Kondorosy E (2021) Taxonomic position of *Heissothignus* and review of the species of *Meschia* (Hemiptera: Heteroptera: Meschiidae). Zootaxa 5032(2): 283–289. https://doi.org/10.11646/zootaxa.5032.2.9

Mancini C (1939) Hemiptera. In: Missione Biologica nel paese dei Borana. Raccolte zoologiche. Reale Accademia d'Italia, Roma. 195–224.

Mancini C (1954) Contributo alla conoscenza degli Emitteri dell'Eritrea. Atti del Museo Civico di Storia Naturale di Trieste 19(3): 137–159.

Mancini C (1956) Emitteri dell'Abissinia raccolti dal M.se Saverio Patrizi e da altri. Nota I. Fragmenta Entomologica 2(8): 65–96.

Masłowski A, Baňař P, Carapezza A, Chérot F, Jindra Z, Taszakowski A, Wolski A (2023) Notes on plant bugs (Hemiptera, Heteroptera, Miridae) from Madagascar with a description of new species of the cylapine tribe Fulviini and checklist of Madagascan mirids. Zootaxa 5330(1): 73–92. https://doi.org/10.11646/zootaxa.5330.1.3

Matocq A (2021) Description de deux nouvelles especes de *Collaria* Provancher de Madagascar (Hemiptera, Miridae, Mirinae, Stenodemini). Bulletin de la Société Entomologique de France 126(4): 429–436. https://doi.org/10.32475/bsef_2183

Montandon AL (1899) III. Hémiptères. In: Junod HA (ed.): La faune entomologique du Delagoa. Bulletin de la Société Vaudoise des Sciences Naturelles 35: 216–220.

Nicholas V, Carapezza A, Rider DA, Kment P (2024) New records, diagnostics and preliminary checklist of the superfamily Pentatomoidea (Hemiptera: Heteroptera) from the Comoro Islands. Zootaxa 5481(1): 1–29. https://doi.org/10.11646/zootaxa.5481.1.1

O'Donnell JE, Schaefer CW (2011) *Lilliputocoris* (Hemiptera: Heteroptera: Rhyparochromidae: Rhyparochrominae: Lilliputocorini) now known from Madagascar. Heteropterus Revista de Entomología 11: 311–316.

Paulian R, Blanc Ch, Guillaumet J-L, Betsch J-M, Griveaud P, Peyrieras A (1973) Étude des écosystemès montagnards dans la région malgache. II. Les chaines Anosyennes. Géomorphologie, climatologie et groupements végétaux. (Campagne RCP 225, 1971–1972). Bulletin du Muséum National d'Histoire Naturelle, 3e Série 118 (Écologie Générale Vol. 1): 1–40.

Pluot-Sigwalt D, Lis JA (2008) Morphology of the spermatheca in the Cydnidae (Hemiptera: Heteroptera): bearing of its diversity on classification and phylogeny. European Journal of Entomology 105(2): 279–312. https://doi.org/10.14411/eje.2008.038

Ralimanana H, Perrigo AL, Smith RJ, Borrell JS, Faurby S, Rajaonah MT, Randriamboavonjy T, Vorontsova MS, Cooke RSC, Phelps LN, Sayol F, Andela N, Andermann T, Andriamanohera AM, Andriambololonera S, Bachman SP, Bacon CD, Baker WJ, Belluardo F, Birkinshaw C, Cable S, Canales NA, Carrillo JD, Clegg R, Clubbe C, Crottini A, Damasco G, Dhanda S, Edler D, Farooq H, Ferreira P de L, Fisher BL, Forest F, Gardiner LM, Goodman SM, Grace OM, Guedes TB, Hackel J, Henniges MC, Hill R, Lehmann CER, Lowry PP, Marline L, Matos-Maraví P, Moat J, Neves B, Nogueira MGC, Onstein RE, Papadopulos AST, Perez-Escobar OA, Phillipson PB, Pironon S, Przelomska NAS, Rabarimanarivo M, Rabehevitra D, Raharimampionona J, Rajaonary F, Rajaovelona LR, Rakotoarinivo M, Rakotoarisoa AA, Rakotoarisoa SE, Rakotomalala HN, Rakotonasolo F, Ralaiveloarisoa BA, Ramirez-Herranz M, Randriamamonjy JEN, Randrianasolo V, Rasolohery A, Ratsifandrihamanana AN, Ravololomanana N, Razafiniary V, Razanajatovo H, Razanatsoa E (2022) Madagascar's extraordinary biodiversity: Threats and opportunities. Science 378: 6623. https://doi.org/10.1126/science.adf1466

Rédei D, Tsai J-F (2021) *Zhengica cornuta* gen. nov., sp. nov., a new genus and species of Pentatomidae from East Asia (Hemiptera: Heteroptera). Zootaxa 5071(2): 271–282. https://doi.org/10.11646/zootaxa.5071.2.6

Rédei D, Tsai J-F (2022) A revision of *Scutellera* (Hemiptera: Heteroptera: Scutelleridae). Zootaxa 5092(1): 1–40. https://doi.org/10.11646/zootaxa.5092.1.1

Reiche L, Fairmaire L (1849) Ordre des Hémiptères. In: Ferret A, Galinier (Ed.) Voyage en Abyssinie 1839–1845. Vol. 3. Paulin Libraire, Paris, 433–450 pp.

Rider DA, Schwertner CF, Vilímová J, Rédei D, Kment P, Thomas DB (2018) Higher systematics of Pentatomoidea. Pp. 25–201. In: McPherson JE (Ed.) Invasive stink bugs and related species (Pentatomoidea): Biology, higher systematics, semiochemistry, and management. CRC Press, Taylor and Francis Group, Boca Raton, London, New York, xix + 819 pp. https://doi.org/10.1201/9781315371221-2

Robertson IAD (2009) The Pentatomoidea (Hemiptera: Heteroptera) of Sub-Saharan Africa. A database. African Studies Centre, Leiden, Netherlands, 451 pp. https://repository.naturalis.nl/pub/391284/Robertson_Pentatomoidea_Africa_2009.pdf

Roca-Cusachs M, Kment P (2022) *Joppeicus paradoxus* (Hemiptera: Heteroptera: Joppeicidae): a new alien species in the European Union? Zootaxa 5195(3): 256–266. https://doi.org/10.11646/zootaxa.5195.3.4

Roca-Cusachs M, Schwertner CF, Kim J-G, Eger J, Grazia J, Jung S-H (2022) Opening Pandora's box: molecular phylogeny of the stink bugs (Hemiptera: Heteroptera: Pentatomidae) reveals great incongruences in the current classification. Systematic Entomology 47(1): 36–51. https://doi.org/10.1111/syen.12514

Roca-Cusachs M, Rider DA, Kment P, Jung S-H (2024) Revision of *Vitruvius* (Hemiptera: Heteroptera: Pentatomidae) with the description of a new species from Cambodia and Thailand. Zootaxa 5486(3): 419–434. https://doi.org/10.11646/zootaxa.5486.3.5

Salini S, Kment P, Cassis G (2023) Rehabilitation of *Mormoschema* (Hemiptera: Heteroptera: Pentatomidae): Generic status restituted and tribal placement revised. Acta Entomologica Musei Nationalis Pragae 63(2): 397–412. https://doi.org/10.37520/aemnp.2023.025

Salini S, Kment P, Lemaître VA, Webb MD, Roca-Cusachs M (2025) Revision of the Austro-Oriental shield-bug genus *Axiagastus* (Hemiptera: Heteroptera: Pentatomidae), with the description of five new species, and taxonomic notes on related genera. Zootaxa 5603(1): 1–82. https://doi.org/10.11646/zootaxa.5603.1.1

Schaefer CW (1981) The sound-producing structures of some primitive Pentatomoidea (Hemiptera: Heteroptera). Journal of the New York Entomological Society 88: 230–235.

Schaefer CW, Dolling WR, Tachikawa S (1988) The shieldbug genus *Parastrachia* and its position within the Pentatomoidea (Insecta: Hemiptera). Zoological Journal of the Linnean Society 93(4): 283–311. https://doi.org/10.1111/j.1096-3642.1988.tb01365.x

Schouteden H (1905) Expédition du Baron C. von Erlanger en Abyssinie et au pays des Somalis. Hémiptères-Hétéroptères. Annales de la Société Entomologique de Belgique 49: 11–16.

Schouteden H (1910) 12. Hemiptera. 6. Pentatomidae. Pp. 73–96. In: Wissenschaftliche Ergebnisse der Schwedischen zoologischen Expedition nach dem Kilimandjaro, dem Meru und den Umgebenden Massaisteppen Deutsch-Ostafrikas 1905–1906 unter Leitung von Prof. Dr. Yngve Sjöstedt. herausgegeben mit Unterstützung von der Königl. Schwedischen Akademie der Wissenschaften. Band 2. Schwedischen Akademie der Wissenschaften, Stockholm, 844 pp. [+ 19 pls]

Schouteden H (1913) Cimicidæ et Coreidæ recueillis au Congo par le Dr. J. Bequaert. Revue Zoologique Africaine 2(2): 189–202.

Schouteden H (1957) Contributions à l'étude de la faune entomologique du Ruanda-Urundi (Mission P. Basilewsky 1953). CXXVI. Heteroptera Plataspidae, Acanthosomidae [sic!], Pentatomidae et Coreidae. Annales du Musée Royal du Congo Belge. Sciences Zoologiques 58: 269–310.

Schuh RT, Slater JA (1995) True bugs of the World (Hemiptera: Heteroptera). Classification and natural history. Cornell University Press, Ithaca and London, 337 pp.

Schumacher F (1913) Ein Beitrag zur Kenntnis der Rhynchoten-Fauna Südafrikas, insbesondere von Deutsch-Südwestafrika, Klein-Namaland und dem Kalaharigebiet. In: Zoologische und Anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Südafrika ausgeführt in den Jahren 1903–1905 mit Unterstützung der Kgl. Preussischen Akademie der Wissenschaften zu Berlin von Dr. Leonhard Schultze Professor der Geographie an der Universität Kiel. Fünfter Band. Systematik und Tiergeographie. Zweite Lieferung. Denkschriften der Medicinisch-Naturwissenschaftlichen Gesellschaft, 17: 49–88.

Signoret V (1880) Deux descriptions et des observations sur divers Hémiptères. Bulletin de la Société Entomologique de France 1880: 193–194.

Signoret V (1881) Révision du Groupe des Cydnides de la Famille des Pentatomides. Partie 1–4. Annales de la Société Entomologique de France Série 6 1: 25–52, 193–218, 319–332, 423–436.

Signoret V (1882) Révision du Groupe des Cydnides de la Famille des Pentatomides. Partie 5–8. Annales de la Société Entomologique de France, Série 6 2: 23–42, 145–168, 241–266, 465–484.

Signoret V (1883) Révision du Groupe des Cydnides de la Famille des Pentatomides. Partie 9–12. Annales de la Société Entomologique de France, Série 6 3: 33–60, 207–220, 357–374, 517–534.

Signoret V (1884) Révision du Groupe des Cydnides de la Famille des Pentatomides. 13e et derniére partie. Annales de la Société Entomologique de France, Série 6 4: 45–62. https://doi.org/10.5962/bhl.title.13250

Sites R, Bergsten J (2022) The Naucoridae (Heteroptera: Nepomorpha) of Madagascar, with revisions of Temnocoris and Tsingala (Laccocorinae). PLoS ONE 17(9) (e0272965): 1–79.. https://doi.org/10.1371/journal.pone.0272965

Stål C (1865) Hemiptera Africana. Tomus I. Officina Norstedtiana, Holmiae, [1864], 256 pp.

Stål C (1866) Hemiptera Africana. Tomus IV. Officina Norstedtiana, Holmiae, 181 pp.

Stål C (1876) Enumeratio Hemipterorum. Bidrag till en förteckning öfver alla hittills kända Hemiptera, jemte systematiska meddelanden. 5. [Checklist of Hemiptera. Contribution to a list of all hitherto known Hemiptera, along with systematic notes. 5]. Kongliga Svenska Vetenskaps-Akademiens Handlingar 14(4): 1–162. [in Latin]

Sweet MH, Schaefer CW (2002) Parastrachiinae (Hemiptera: Cydnidae) raised to family level. Annals of the Entomological Society of America 95(4): 441–448. https://doi.org/10.1603/0013-8746(2002)095[0441:PHCRTF]2.0.CO;2

Taszakowski A, Masłowski A, Wolski A, Gorczyca J (2022) Two new genera and species of the Fulviini (Hemiptera, Heteroptera, Miridae, Cylapinae) from Madagascar, with a checklist of Madagascan cylapines. Zootaxa 5209(1): 93–110. https://doi.org/10.11646/zootaxa.5209.1.5

Taszakowski A, Jindra Z, Wolski A (2025a) *Gryllofulvius gibbosus* Taszakowski gen. et sp. nov. – a remarkable, flightless and stridulating plant bug (Heteroptera: Miridae: Cylapinae) from Madagascar. European Journal of Taxonomy 976: 171–181. https://doi.org/10.5852/ejt.2025.976.2785

Taszakowski A, Kim J-G, Kaszyca-Taszakowska N, Gierlasiński G, Jung S-H (2025b) Taxonomic review of the genus *Psallops* Usinger, 1946 (Hemiptera: Miridae) from the Afrotropical Region, with descriptions of two new species and a discussion on the internal revision of *Psallops*. Zootaxa 5646(4): 545–561. https://doi.org/10.11646/zootaxa.5646.4.4

Thunberg CP (1783) Dissertatio entomologica novas insectorum species, sistens, cujus partem secundam, cons. exper. facult. med. upsal., publice ventilandam exhibent. Johan. Edman, Upsaliae, 29–52. [pl. 2]

Tsai J-F, Rédei D, Yeh G-F, Yang M-M (2011) Jewel bugs of Taiwan (Heteroptera: Scutelleridae). National Chung Hsing University, Taichung, 309 pp.

Villiers A (1952) La réserve naturelle intégrale du Mt. Nimba. Fascicule 1. XIV. Hémiptères Hétéroptères terrestres. Mémoires de l'Institut Français d'Afrique Noire 19: 289–309.

Walker F (1867a) Catalogue of the specimens of Heteropterous Hemiptera in the collection of the British Museum. Part II. Scutata. Trustees of the British Museum, London, 1–240. https://doi.org/10.5962/bhl.title.118688

Walker F (1867b) Catalogue of the specimens of Heteropterous Hemiptera in the collection of the British Museum. Part II. Scutata. Trustees of the British Museum, London, 241–417. https://doi.org/10.5962/bhl.title.118688

Wallengren HDJ (1875) Insecta Transvaaliensia. – Bidrag till Transvaalska Republikens i Södra Afrika Insektfauna. [Insects of Transvaal. – Contribution to the insect fauna of the Transvaal Republic in South Africa]. Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar 32(1): 83–137. [in Swedish and Latin]

Westwood JO (1837) [Descriptions.] In: Hope FW: A Catalogue of Hemiptera in the Collection of the Rev. F. W. Hope, M. A. with short Latin diagnoses of the new species. Part 1. London, 46 pp. [Part 2, pp. 1–26 (1842) states that descriptions are by J. O. Westwood.]

Wu Y-Z, Yu Sh-Sh, Wang Y-H, Wu H-Y, Li X-R, Men X-Y, Zhang Y-W, Rédei D, Xie Q, Bu W-J (2016) The evolutionary position of Lestoniidae revealed by molecular autapomorphies in the secondary structure of rRNA besides phylogenetic reconstruction (Insecta: Hemiptera: Heteroptera). Zoological Journal of the Linnean Society 177(4): 750–763. https://doi.org/10.1111/zoj.12385

Zámbó A, Baňař P, Kovács S, Kondorosy E (2022) Drymini of Madagascar, with description of a new genus and three new species (Hemiptera: Heteroptera: Rhyparochromidae). Acta Entomologica Musei Nationalis Pragae 62(1): 211–223. https://doi.org/10.37520/aemnp.2022.016

Zettel H (2020) New water strider species of *Eurymetra* from Madagascar (Hemiptera: Heteroptera: Gerridae). Acta Entomologica Musei Nationalis Pragae 60(1): 15–22. https://doi.org/10.37520/aemnp.2020.002

Zettel H, Laciny A (2020) *Rhagovelia* species from Nosy Boraha east of Madagascar (Hemiptera: Heteroptera: Veliidae). Acta Entomologica Musei Nationalis Pragae 60(2): 419–425. https://doi.org/10.37520/aemnp.2020.26

Zhou Y-Y, Rédei D (2018) A new synonymy in East Asian Urostylididae (Hemiptera: Heteroptera). Zootaxa 4504(1): 145–150. https://doi.org/10.11646/zootaxa.4504.1.10

Zhou Y-Y, Rédei D (2020) From lanceolate to plate-like: Gross morphology, terminology, and evolutionary trends of the trichophoran ovipositor. Arthropod Structure & Development 54(100914): 1–29. https://doi.org/10.1016/j.asd.2020.100914

Zrzavý J (1990) Evolution in antennal sclerites in Heteroptera (Insecta). Acta Universitatis Carolinae. Biologica 34: 189–227.